

Clinical Cancer Research

Peroxisome Proliferator-Activated Receptor γ Pathway Targeting in Carcinogenesis: Implications for Chemoprevention

Frank Ondrey

Clin Cancer Res 2009;15:2-8. Published online December 31, 2008.

Updated Version Access the most recent version of this article at:
doi:[10.1158/1078-0432.CCR-08-0326](https://doi.org/10.1158/1078-0432.CCR-08-0326)

Cited Articles This article cites 72 articles, 37 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/15/1/2.full.html#ref-list-1>

Citing Articles This article has been cited by 6 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/15/1/2.full.html#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.

Peroxisome Proliferator-Activated Receptor γ Pathway Targeting in Carcinogenesis: Implications for Chemoprevention

Frank Ondrey

Abstract The peroxisome proliferator-activated receptor (PPAR) γ is one member of the nuclear receptor superfamily that contains in excess of 80 described receptors. PPAR γ activators are a diverse group of agents that range from endogenous fatty acids or derivatives (linolenic, linoleic, and 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂) to Food and Drug Administration-approved thiazolidinedione drugs [pioglitazone (Actos) and rosiglitazone (Avandia)] for the treatment of diabetes. Once activated, PPAR γ will preferentially bind with retinoid X receptor α and signal antiproliferative, antiangiogenic, and prodifferentiation pathways in several tissue types, thus making it a highly useful target for down-regulation of carcinogenesis. Although PPAR- γ activators show many anticancer effects on cell lines, their advancement into human advanced cancer clinical trials has met with limited success. This article will review translational findings in PPAR γ activation and targeting in carcinogenesis prevention as they relate to the potential use of PPAR γ activators clinically as cancer chemoprevention strategies.

Over the past 20 years, strides have been made in treatment of several solid tumor malignancies resulting in measurable increases in survival, but a firm challenge remains to establish improved cancer prevention strategies for most cancers. Strategies focusing on abrogating global events during human carcinogenesis [e.g., epigenetic events with histone deacetylase (HDAC) inhibitors, carcinogenesis with natural products, or drugs affecting transcription factor pathways] coexist with focused strategies that seem highly specific such as individual receptor inhibition (e.g., epidermal growth factor receptor) or single-enzyme inhibition (e.g., cyclooxygenase). Presently, antiestrogen strategies for breast carcinoma, topical agents such as Aldara and Effudex for skin carcinoma, and cyclooxygenase-2 inhibitors (celecoxib) for familial polyposis are some of a very limited number of Food and Drug Administration-approved strategies for chemoprevention. Other promising therapies (e.g., retinoids and cyclooxygenase-2 targeting) have fallen from favor due to unacceptable toxicity (1, 2) or inefficacy (3, 4). There is a demand for safe agents that target high-risk conditions such as preexisting intraepithelial neoplasia, a high-risk cancer precursor (ref. 5; Table 1). Peroxisome proliferator-activated receptor γ (PPAR- γ) activation strategies may help fulfill this demand.

Nuclear Receptor Peroxisome Proliferator-Activated Receptor Activators Broadly Target Preneoplastic Processes

The successful exploitation of nuclear receptor targeting for cancer prevention have thus far included antiestrogen strategies for breast carcinoma prevention. In this scenario, the estrogen receptor, which is linked to breast cancer growth, is successfully antagonized by tamoxifen or raloxifene. Conversely, the agonism of other nuclear receptors (e.g., PPARs) that are more often associated with differentiation is another strategy to target this receptor class.

Peroxisome proliferator-activated receptors α , β (or δ), and γ are 3 of ~100 nuclear receptors in the orphan receptor class (6, 7). This receptor class consists of a variety of steroid, retinoid, thyroid, vitamin D, and other receptors that dimerize and regulate a multitude of downstream metabolic processes when activated or inhibited. The original discovery and cloning of the PPARs leads to a somewhat confusing explanation of their functions because of their tissue distribution and differential effects in humans versus animal models and the fact that they diffusely influence basic metabolic processes, such as energy storage. For example, PPAR α is the downstream receptor for a variety of peroxisome proliferators (including industrial plasticizers) associated with hepatocellular carcinogenesis in rats but not in humans (8, 9). Conversely, in humans, natural ligands for PPAR α include a variety of fatty acids responsible for lipid storage and are clearly not carcinogens. Synthetic agents, such as fibrates, are approved for the treatment of hyperlipidemia, target PPAR α , and are also not carcinogenic in humans. However, at high doses, combined agonists of the PPAR α /PPAR γ class are associated with bladder carcinogenesis in animals and this has affected their utility clinically (10). The complex nature of these signaling pathways in different species creates these paradoxes.

PPAR γ was originally described as a differentiation transcription factor for adipose tissue (11). Its natural ligands are felt to

Authors' Affiliation: Department of Otolaryngology and University of Minnesota Masonic Cancer Center, University of Minnesota, Minneapolis, Minnesota
Received 4/23/08; revised 8/3/08; accepted 8/11/08.

Requests for reprints: Frank Ondrey, Department of Otolaryngology and University of Minnesota Masonic Cancer Center, University of Minnesota, 410 Delaware Street Southeast, MMC396, Minneapolis, MN 55455. Phone: 612-625-9449; Fax: 612-625-2101; E-mail: ondre002@umn.edu.

©2009 American Association for Cancer Research.
doi:10.1158/1078-0432.CCR-08-0326

Table 1. Drugs affecting PPAR γ signaling

Agent	Receptor	Drug class	Food and Drug Administration approved
Pioglitazone	PPAR γ	Thiazolidinedione	Yes
Rosiglitazone	PPAR γ	Thiazolidinedione	Yes
Ciglitazone	PPAR γ	Thiazolidinedione	No
Troglitazone	PPAR γ	Thiazolidinedione	Yes/withdrawn
Farglitazar	PPAR γ	Thiazolidinedione	No
Muraglitazar	PPAR α,γ	Thiazolidinedione	No
GW7845	PPAR γ	Unknown	No
GW7875	PPAR γ	Unknown	No
GW501516	PPAR δ	Unknown	No
LG10068	RXR α	Retinoid	No
Clofibrate	PPAR α	Fibrate	Yes
Prostaglandin J ₂	PPAR γ	Prostaglandin	No

be products of the eicosanoid cascade and 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ has a particular affinity for this receptor. The thiazolidinedione drugs are used for the treatment of type II diabetes and specifically target PPAR γ (Table 1). PPAR δ has been associated with colon carcinogenesis in some models and was originally felt to not be particularly related to any logical metabolic processes. Recently, a broader understanding of these receptors has been placed into context in an atlas of nuclear receptors whereby the PPARs are understood as being principally associated with nutrient uptake, storage, and utilization during the circadian rhythm.^{1,2} Therefore, an elevated importance is ascribed to these receptors and the utility of targeting them in conditions such as obesity, diabetes, and other metabolic processes such as exercise is emerging. Currently, Food and Drug Administration-approved agents are available to target PPAR α and γ .

Preferential binding partners for PPAR- γ include receptors of the retinoic acid class. The retinoic acid receptors consist of two families and six members: the retinoic acid receptors (α , β , and γ) and the retinoid X receptor (RXR; α , β , and γ). By themselves, this class of receptors can heterodimerize, but in the past several years, a transcriptome consisting of RXR/retinoic acid receptor heterodimers is linked to HDAC has been identified (12–14). This is derepressed on activation with 9-*cis* retinoic acid binding to RXR elements. There are a large number of natural and synthetic retinoids that target these receptors, and some retinoid ligands can bind to either class of receptors (e.g., all-*trans* retinoic acid), whereas others are class restricted (e.g., 9-*cis* retinoic acid and the RXR family). Dysfunction within this signaling complex may therefore be targeted with a large number of drugs affecting either retinoid or HDAC receptors to down-regulate carcinogenesis. This axis, which putatively also involves PPARs, could therefore be dually targeted with drugs that affect retinoid or HDAC receptors, or PPAR and HDAC receptors, once the key dysregulated points are identified. Additionally, simply the presence of functional receptors would allow for exploitation within this axis. One example would be the modulation of receptors of the retinoic acid receptor β class in aerodigestive chemoprevention, which has been shown to be up-regulated in successful aerodigestive chemoprevention (15–18).

¹Margolis RN, Evans RM, O'Malley BW, NURSA Atlas Consortium. The Nuclear Receptor Signaling Atlas: development of a functional atlas of nuclear receptors. *Mol Endocrinol* 2005;19:2433–6.

²Yang X, Downes M, Yu RT, et al. Nuclear receptor expression links the circadian clock to metabolism. *Cell* 2006;126:801–10.

Molecularly, it is understood that PPAR γ will form a heterodimer with RXR α and transcriptionally activate downstream genes (Fig. 1; ref. 19). The PPAR γ /RXR α complex can be phosphorylated at either protein (20–22) and the transcription factor will preferentially bind direct repeat sequences in downstream genes. The consequences of phosphorylation by mitogen-activated protein kinase or other factors (e.g., insulin) is not well studied but in general leads to attenuation of receptor-dependent and receptor-independent effects, such as attenuation of adipose lineage genes in preadipocytes. Interactions occurring in the transcription factor complex with other cofactors (such as SMRT and NCoR), which further modulate PPAR γ activity, have been best described in adipogenesis models (23–26).

In cancer, the interactions between PPAR γ /RXR α have been the principal targets for exploitation compared with the study of metabolic diseases or adipose metabolism and obesity. The drugs utilized for cancer therapy studies (that would also be appropriate for chemoprevention studies) have included the preclinical activators of PPAR (such as prostaglandins of the J series) and dietary additions (such as linoleic and linolenic acid derivatives). Thiazolidinediones available either preclinically (such as ciglitazone) or as Food and Drug Administration approved drugs for the treatment of diabetes (such as pioglitazone and rosiglitazone) have been used. Additionally, a large number of natural and synthetic retinoid receptor targeting agents have been studied with the focus on RXR/retinoic acid receptor interactions in cancer. There has been less focus on molecular events occurring along the PPAR- γ /RXR- α axis. Further, the more recently described interactions of the RXR with HDAC could additionally be studied with the focus of PPAR γ and downstream events. Finally, drugs affecting PPAR γ or RXR- α phosphorylation may also be useful (MAPK inhibitors).

Clinical-Translational Advances

PPARs and cancer treatment. The literature is replete with a large volume of studies showing efficacy of PPAR treatment of cell lines and animal models. It is well established that activating each subclass of receptors can result in antiproliferative, proapoptotic, prodifferentiation, and antiangiogenic effects in cell lines or animal models (27). These data, collectively, have resulted in several completed clinical trials in solid tumor malignancies in breast (28), prostate (29), and liposarcoma

(30). Unfortunately, there is a lack of clinical effect in advanced refractory malignancies when PPAR γ agonists are used as targeted agents alone. Because the agents available are pleiotropic in nature and given at therapeutic and pharmacologic doses in publications, their effects in a given target organ may vary; both PPAR γ -dependent and PPAR γ -independent effects could be occurring simultaneously in the same tissue or cell line. This makes it difficult to form general consensus statements regarding their effects in human cancers. In light of clinical inefficacy of the agents, perhaps these agents should be thought of as reasonably safe adjuncts to standard chemotherapy in further testing.

Organ site-specific carcinogenesis and chemoprevention studies of PPAR γ . Because the original discovery of the PPAR activators was potentially associated with carcinogenesis, careful evaluation of these agents and metabolic pathways is required to delineate doses and conditions whereby untoward effects may occur (8, 31). Several studies have investigated PPAR γ as therapy for carcinogenesis prevention preclinically or preneoplasia, with other support for specific mechanisms of action in target organs (e.g., receptor mutations, allelic variation, etc.).

Thyroid. In human thyroid follicular carcinoma, PPAR γ can form a fusion oncogene with the PAX8 promoter (32). This fusion oncogene has been described in several studies and confirmed by numerous techniques and is detectable in up to 50% of follicular thyroid carcinomas. Additionally, it has also been discovered in thyroid follicular adenomas (33), which are typically surgically removed as carcinoma precursors. The discovery of this fusion oncogene is potentially concerning but multiple studies have shown both increased and more differentiated phenotypes of the tumors, which seem to be associated with the t(2;3)(q13;p25) translocation responsible for the PAX8/PPAR γ protein. It has been established that this fusion oncoprotein can activate the PPAR response element

when transfected into Rat FRTL-5 thyroid cells and also bind the PPAR response element on electromobility shift assay and this resulted in overall increased cell proliferation (34). However, in this analysis, some PPAR γ -responsive genes were activated by the protein and others were inhibited. Therefore, this mutation can act by both gain and loss of function in thyroid carcinoma, the balance of which favors differentiated tumor formation in affected individuals. It is possible in instances where this oncoprotein is associated with more aggressive tumors, the fusion gene may possess a greater degree of gain of function attributable to transcription factor and tumor-specific PPAR γ cofactor function and regulation. In contradistinction, in the TR β ^{PV/PV} mouse (which contains a mutated thyroid hormone receptor and develops follicular neoplasms spontaneously), the TR β receptor acts as a dominant-negative on normal PPAR γ transactivation (35). Currently, follicular thyroid neoplasms are the only known neoplasms to be associated with PPAR γ fusion gene products and PAX 8/PPAR γ is the only known PPAR γ -associated fusion gene.

Gastrointestinal system. Earlier studies in colon carcinogenesis demonstrated a potential role for PPAR γ in colorectal tumor growth. On PPAR δ principally based on early work that suggested a role for PPAR δ in colorectal tumor growth (36, 37). In colon carcinoma, experimental evidence is somewhat ambiguous in that PPAR γ may play a protective or tumor-suppressive role against cancer development as well as tumorigenesis. Original experiments in the Min mouse model showed enhanced noninvasive polyp formation in the colons of mice given pharmacologic concentrations of troglitazone in the diet for 7 weeks (38). Histologically, the polyps showed dysplasia and hyperplasia without invasion. No alterations were observed in the small intestinal rate of polyp formation with troglitazone. Lefebvre et al. performed a second set of experiments with troglitazone and a second thiazolidinedione, BRL49,653, with similar findings in both the large and the




Fig. 1. Evidence-based PPAR γ targeting in carcinogenesis. PPAR γ will form a heterodimer with RXR α . It is possible that the RXR α can be derepressed from an association with HDAC to facilitate this new binding. Once associated, the PPAR γ may be activated with a thiazolidinedione drug to become an "activated" heterodimer, which, in the nucleus, can trigger several downstream gene activation events. These include decreases in proliferation through either cyclin D1 or cyclin-dependent kinases 1, 2, and 4 as well as Ki-67 and proliferating cell nuclear antigen. Other families of either adipose or tissue lineage-specific differentiation proteins are up-regulated, contributing to a more differentiated cell. Also, genes contributing to events, such as CD31 and angiogenesis, may be down-regulated by the activation of PPAR γ . Apoptosis-associated proteins such as proline oxidase can also be activated by PPAR γ . Finally, PPAR γ can target further expression of itself, creating a positive feedback loop, thus amplifying the original effect of the drug-mediated targeting of the receptor.

small bowel (39). Similarly, Yang et al. described troglitazone-induced colon tumorigenesis when given at 0.2% in the diet in C57BL/6J normal mice as well as in *Apc*^{L638N/+} *Mlh1*^{+/-} mice (40). More recently, Linsalata et al. found enhanced expression of PPAR γ and ornithine decarboxylase pathway products in human tumors compared with normal mucosa, and this was more prevalent in colon cancers with ras mutations (41). These four studies support a role for PPAR γ potentially playing a role in colon carcinogenesis albeit primarily in transgenic mouse models.

Conversely, other studies have provided evidence for PPAR γ activation as a way to abrogate colon carcinogenesis. Ulcerative colitis is a disease associated with chronic inflammation and increased large bowel cancers. It has recently been discovered that the anti-inflammatory agent 5-aminosalicylic acid, used to treat ulcerative colitis, stimulates PPAR γ and is anticarcinogenic. Ulcerative colitis-associated colon cancers are modeled with oral dextran sulfate/intraperitoneal azoxymethane (42). With 100 mg/kg/d 5-aminosalicylic acid administration for 7 weeks, tumorigenesis was decreased by 50% and the tumors were statistically smaller. With 25 mg/kg (low dose) administration of pioglitazone or rosiglitazone, there was no decrease in tumorigenesis over controls. Others have shown that PPAR γ mRNA is induced by 5-aminosalicylic acid treatment in humans and PPAR γ is deficient in patients afflicted with ulcerative colitis (43). Another potentially useful chemopreventive therapeutic, 9,11-conjugated linoleic acid, has also been tested as a potential PPAR γ -activating treatment for the prevention of colon carcinogenesis. This agent is derived from bitter melons, and when administered in the diet of azoxymethane-treated Fischer rats, 50% decreases in tumor incidence and size were observed in the 9,11-conjugated linoleic acid-treated rats, and this was accompanied by an increase in the level of PPAR γ in colonic mucosa (44). Others have shown PPAR γ activation with 9,11-conjugated linoleic acid (45, 46), and because PPAR γ is known to induce itself, this is a simple explanation for why elevated PPAR γ levels were observed. In another study that supports an anticarcinogenic role for PPAR γ in colon carcinogenesis, immortalized rat intestinal IEC6 cells experience reduced proliferation with linoleic acid (47). These changes are reversible when PPAR γ expression is inhibited by RNA knockdown techniques, thus suggesting a tumor-suppressive role for PPAR γ constitutive expression. Finally, it has recently been pointed out that thiazolidinedione treatment enhances apoptosis in colon cancer cells through the activation of proline oxidase (48). In other gastrointestinal system malignancies, thiazolidinedione administration with RXR agonists in rat hepatocyte cell lines induces chemopreventive glutathione S-transferase 2 α (49). In *N*-methyl-*N*-nitrosourea-induced gastric carcinoma, PPAR γ ^{+/-} transgenic mice developed fewer tumors than their heterozygous (PPAR γ ^{+/-}) counterparts, but troglitazone was only effective in preventing tumors in the homozygotes (50). These data, taken together, show that troglitazone, given at pharmacologic doses, can promote tumor formation in the Min mouse model and in one study using wild-type mice. This agent is no longer on the market. However, other studies show that the induction of PPAR γ may be protective against the development of these colon and other gastrointestinal malignancies and promote apoptosis. Finally, agents such as 5-aminosalicylic acid and 9,11-conjugated linoleic acid may

be chemopreventive for colon carcinogenesis. Further studies with FDA-approved PPAR γ activators and intestinal carcinogenesis are lacking.

Breast. The evidence for utility of PPAR γ activators in the prevention of breast cancer is perhaps more straightforward than observed in the conflicting reports in colon cancer development. Two studies (one in mice and one in rats) conducted in the late 1990s showed efficacy in the prevention of mammary carcinogenesis with troglitazone and GW7845. Mehta et al. (51) employed a BALB/c 7,12-dimethylbenz[*a*]anthracene organ culture, which showed a 50% to 100% decrease in gland lesions after treatment with troglitazone or troglitazone and the RXR-selective agonist, LG10068. In this model, the best treatment was a combination of troglitazone and LG10068. Suh et al. (52) used GW7845 in a rat *N*-methyl-*N*-nitrosourea model of breast carcinogenesis and identified decreases in tumor incidence, tumors per animal, and tumor weight after 2 months of treatment. More recently, both PPAR γ and PPAR δ activators were used to study 7,12-dimethylbenz[*a*]anthracene-treated FVB mice (53). Interestingly, the PPAR γ activator GW7875 inhibited tumorigenesis; the tumors that formed were ductal in origin and expressed estrogen receptor- α . The PPAR δ activator GW501516 actually increased the incidence of tumors over baseline and the histology was either adenocarcinoma or squamous carcinomas. Badawi et al. tested an experimental PPAR γ activator, Fmoc-L-Leu and celecoxib, alone or in combination, in a *N*-methyl-*N*-nitrosourea model of rat breast carcinogenesis (54). Both agents were effective in reducing the incidence of tumors by ~30%; however, the combination of both agents was responsible for a 65% decrease in tumor incidence. These treatments separately decreased cyclooxygenase-2 and increased PPAR γ in tissues, and decreases in prostaglandin E₂ were also observed. Apoptosis was increased and measures of proliferation including proliferating cell nuclear antigen and cyclin-dependent kinase 1 were decreased. In a population study of PPAR γ allelic variants isoforms and breast cancer incidence in a Danish population, a variant splice allele of the PPAR γ 2 gene was associated with decreased risk in women identified as alcohol users (55). These studies point out class differences in the effect of PPAR activators on breast carcinogenesis, with PPAR γ inhibiting tumor formation. The potential for combination chemoprevention of breast carcinoma with PPAR γ and RXR α activation is also shown. Because PPAR activators are not always pure agonists for a specific subclass of receptors, we are to be cautioned that results from experiments of mixed agonists may not allow straightforward interpretation.

Skin. PPAR γ focused studies have been conducted to examine skin squamous malignancies. The original skin carcinogenesis prevention studies occurred in mouse models. He et al. published both troglitazone and ciglitazone decreased proliferation in primary mouse newborn keratinocytes and the CD50 mouse cell line (56). Cell cycle G₁-phase arrest and suppression of cyclin D1 and cyclin-dependent kinases 4 and 2 occurred as well. However, these effects were found to be PPAR γ independent, as transfection with dominant-negative PPAR γ produced no changes in results of troglitazone treatment. This study was followed with an animal study of 7,12-dimethylbenz[*a*]anthracene-induced carcinogenesis where both rosiglitazone and troglitazone had no effect on either 7,12-dimethylbenz[*a*]anthracene/TPA- or UV-induced

carcinogenesis (57). Further, PPAR γ could not be identified in either normal or transformed tissues. However, when PPAR δ is targeted, its activation is accompanied by a protein kinase C mediated down-regulation of kinases, which ultimately down-regulate extracellular signal-regulated kinase (58). These studies do not support a hypothesis-driven role for PPAR γ activator use in the prevention of skin carcinogenesis.

Aerodigestive carcinogenesis. In aerodigestive squamous carcinoma, the scenario is different than in squamous cancer. In the rat 4-nitroquinolone model of oral carcinogenesis showed ~40% decreases in the incidence of oral squamous carcinoma with decreased multiplicity in rats treated with 500 ppm pioglitazone (59) was observed. Forty-percent reductions in the incidence of rat tongue carcinomas were also observed in a separate study with troglitazone (60). Further support of these preclinical findings can be derived from studies of Govindarajan et al. (61), who were able to show decreased incidence of both head and neck (>40%) and lung (>30%) neoplasms in a Veterans' Administration population of diabetics aged >40 years who were taking thiazolidinedione agents for at least a year. Preclinical studies in lung cancer treatment with PPAR γ agonists were done by Chang and Szabo (62), who showed that proliferative indices were decreased and apoptosis and differentiation markers were increased in non-small cell lung cancer cell lines treated with ciglitazone. These data show that skin and oral squamous carcinoma models respond differently to treatment with PPAR γ activators, with the thiazolidinediones being ineffective in skin squamous carcinogenesis animal models but significantly altering aerodigestive carcinogenesis. The data from lung cancer experiments give further support for hypothesis-driven biomarkers in proliferation, differentiation, and apoptosis for potential chemoprevention studies with these agents. Finally, the combined data from health-care registries of glitazone use combined with animal models provides a stronger rationale to move forward clinically in aerodigestive premalignancy than animal study evidence alone.

Renal system. Another area of investigation has involved the kidney and related organs and PPAR γ effects. The principal experiments in this area have occurred in the bladder urothelial epithelium. In normal urothelial cell cultures, troglitazone is able to induce markers of urothelial differentiation while decreasing squamous metaplasia (63, 64). However, the largest focus on renal epithelial malignancies has been with regard to testing of dual PPAR α/γ agonists in animal models before Food and Drug Administration approval. In this setting, the combined agonists at pharmacologic concentrations result in bladder and/or kidney carcinogenesis with at least five dual agonists (10). One detailed pathologic study provides evidence that chronic effects of urolithiasis contribute to this effect (65). This has resulted in withdrawal of most of these agents from further testing. Interestingly, these studies point out a paradox because high pharmacologic doses in animals result in tumor formation with dual agonists, yet PPAR γ activation promotes a panel of differentiation markers in renal epithelial cells.

Challenge of incorporating PPAR γ agents into prevention clinical trials. One principal issue in any cancer chemoprevention strategy is the safety of the agent. The cardiotoxicity revealed in a recent meta-analysis of rosiglitazone may make this agent less attractive in the prevention realm (66), but pioglitazone does not seem to share this toxicity in a large study of diabetics prospectively analyzed for cardiac risks (67, 68).

Other PPAR γ activators are currently under development but may not gain industry support. Pioglitazone has been used for a phase II cancer prevention trial of preneoplastic aerodigestive lesions³ and could be an important available agent for further testing in intraepithelial neoplasia. Anti-inflammatory agents such as 5-aminosalicylic acid would be other potential agents for PPAR γ -targeted chemoprevention perhaps in high-risk inflammatory bowel disease patients. Also, natural product derivatives such as 9,11-conjugated linoleic acid from bitter melons might be useful in the clinical chemoprevention setting. It is clear that PPAR γ signaling represents a powerful influence in cells where it is metabolically active. It forms a transcriptional complex with RXR α and other cofactors such as NCoR and SMRT. Therefore, combination chemoprevention with agents targeting PPAR γ , RXRs, and related metabolic processes would be reasonable approaches as long as the toxicity of such combinations was acceptable.

Secondly, potential hypothesis-driven surrogate markers in clinical trials emerge from the myriad cell processes affected by PPAR γ activators (Fig. 1). First, it is notable PPAR γ was often induced in tissues as a result of treatment. Therefore, examination of modulation of PPAR γ in tissue samples before and after treatment would be one potentially useful tissue marker of an organ-specific effect in a chemoprevention clinical trial. Additionally, modulation of PPAR γ binding partners (such as RXR α) could also be assayed by immunohistochemistry in tissue specimens before and after PPAR γ activator treatment. The intracellular location or phosphorylation status of the proteins may also be significant, as phosphorylated forms of PPAR γ and RXR α located in the nucleus are probably the most likely isoforms to be found in an active state, thus signifying a tissue-specific treatment effect. Next, several studies examined markers of both proliferation and apoptosis after treatment. Clearly, cyclin D1, proliferating cell nuclear antigen, Ki-67, and cyclin-dependent kinases represent a second class of putative surrogate endpoints, which may undergo change during treatment with PPAR γ activators. Markers of apoptosis including apoptosis index, caspase activation, or proline oxidase induction would also be reasonable biomarkers. Perhaps the most interesting biomarkers might be those associated with cellular differentiation often promoted by PPAR γ activator treatment. These would potentially represent PPAR γ -dependent genes in a variety of target tissues and may provide further insight into nuances of PPAR γ function. They might also represent adipose lineage markers, as several studies indicate that markers of adipose lineage differentiation can occur with PPAR γ treatment (69–71). The antiangiogenic properties represent yet another target for PPAR γ effects and the use of endothelial markers to measure angiogenesis has been advocated (72).

PPARs and current clinical trials. There are several PPAR γ agonists in the pipeline, undergoing clinical trials primarily for metabolic diseases such as diabetes. GSK has at least three novel agents that are PPAR activators in metabolic disease clinical trials (376501, farglitazar, and rosiglitazone XR; GSK product development pipeline, February 2008). One Daiichi Sankyo compound, CS-7017, an oral PPAR γ agonist, is undergoing a single arm phase I evaluation⁴ for advanced metastatic cancer.

³ Unpublished data.

⁴ www.clinicaltrials.gov NCT00408434.

A single-arm combination therapy phase I/II clinical trial employing a taxane and CS-7017, for anaplastic thyroid cancer, is also ongoing (NCT00603941). There are two recently completed single-arm phase IIa leukoplakia reversal clinical trials, one employing rosiglitazone and the other employing pioglitazone.⁵ The pioglitazone clinical trial showed leukoplakia reversal in most patients and a randomized phase II clinical trial with pioglitazone in leukoplakia patients is planned.⁶ These ongoing, or recently completed studies, show considerable ongoing interest in the clinical study of PPAR γ agonists for cancer as well as diabetes.

Future directions. It is clear that the road to safe, effective chemoprevention is arduous and agents in routine use are limited. However, substantial evidence on dosing, potential endpoints in clinical trials, and success in animal models of carcinogenesis provide significant preclinical evidence for how to approach use of these agents in early-phase clinical trials.

⁵ www.clinicaltrials.gov

⁶ Unpublished data.

Thiazolidinediones and natural products discussed that activate PPAR γ may be lead candidate agents for such studies. The risk benefit ratio of the agents needs to be assessed, and this may be further assisted by logical incorporation of biomarkers into the trials. Of great benefit is that many of the biomarkers for proliferation and apoptosis as well as standard pathology from biopsied lesions are now available at most community hospitals and are standardized on automated immune histochemistry. It is easy to become deterred from considering this strategy, particularly in light of the evidence from some transgenic mouse models and the animal studies employing PPAR α/γ combined activators, but it is important to note the antiestrogens commonly in use are associated with endometrial carcinoma development (73). Clearly, risk stratification and the targeting of these agents to specific intraepithelial neoplastic conditions will be important in the future testing of these promising chemoprevention drugs.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

References

- Half EE, Arber N. Chemoprevention of colorectal cancer: two steps forward, one step back? *Future Oncol* 2006;2:697–704.
- Bresalier RS, Sandler RS, Quan H, et al. Cardiovascular events associated with rofecoxib in a colorectal adenoma chemoprevention trial. *N Engl J Med* 2005;352:1092–102.
- Papadimitrakopoulou VA, William WN, Jr., Dannenberg AJ, et al. Pilot randomized phase II study of celecoxib in oral premalignant lesions. *Clin Cancer Res* 2008;14:2095–101.
- Khuri FR, Lee JJ, Lippman SM, et al. Randomized phase III trial of low-dose isotretinoin for prevention of second primary tumors in stage I and II head and neck cancer patients. *J Natl Cancer Inst* 2006;98:441–50.
- O'Shaughnessy JA, Kelloff GJ, Gordon GB, et al. Treatment and prevention of intraepithelial neoplasia: an important target for accelerated new agent development. *Clin Cancer Res* 2002;8:314–46.
- Evans RM. The nuclear receptor superfamily: a Rosetta stone for physiology. *Mol Endocrinol* 2005;19:1429–38.
- Mangelsdorf DJ, Thummel C, Beato M, et al. The nuclear receptor superfamily: the second decade. *Cell* 1995;83:835–9.
- Reddy JK, Lalwai ND. Carcinogenesis by hepatic peroxisome proliferators: evaluation of the risk of hypolipidemic drugs and industrial plasticizers to humans. *Crit Rev Toxicol* 1983;12:1–58.
- Klaunig JE, Babich MA, Baetcke KP, et al. PPAR α agonist-induced rodent tumors: modes of action and human relevance. *Crit Rev Toxicol* 2003;33:655–780.
- NDA 21-865 ATs. Advisory Committee Briefing Document. Preclinical pharmacology and toxicology summary, drug: Pargluva[®] (muraglitazar, BMS-298,585), peroxisome proliferator-activated receptor dual agonist. In: FDA, editor. 2005.
- Kliewer SA, Forman BM, Blumberg B, et al. Differential expression and activation of a family of murine peroxisome proliferator-activated receptors. *Proc Natl Acad Sci U S A* 1994;91:7355–9.
- Altucci L, Gronemeyer H. The promise of retinoids to fight against cancer. *Nat Rev Cancer* 2001;1:181–93.
- Clarke N, Germain P, Altucci L, Gronemeyer H. Retinoids: potential in cancer prevention and therapy. *Expert Rev Mol Med* 2004;6:1–23.
- Okuno M, Kojima S, Matsushima-Nishiwaki R, et al. Retinoids in cancer chemoprevention. *Curr Cancer Drug Targets* 2004;4:285–98.
- Lotan R. Retinoids and their receptors in modulation of differentiation, development, and prevention of head and neck cancers. *Anticancer Res* 1996;16:2415–9.
- Lotan R. Retinoids and chemoprevention of aerodigestive tract cancers. *Cancer Metastasis Rev* 1997;16:349–56.
- Wan H, Dawson MI, Hong WK, Lotan R. Overexpressed activated retinoid X receptors can mediate growth inhibitory effects of retinoids in human carcinoma cells. *J Biol Chem* 1998;273:26915–22.
- Xu XC, Sozzi G, Lee JS, et al. Suppression of retinoic acid receptor β in non-small-cell lung cancer *in vivo*: implications for lung cancer development. *J Natl Cancer Inst* 1997;89:624–9.
- Berger J, Moller DE. The mechanisms of action of PPARs. *Annu Rev Med* 2002;53:409–35.
- Diradourian C, Girard J, Pegorier JP. Phosphorylation of PPARs: from molecular characterization to physiological relevance. *Biochimie* 2005;87:33–8.
- Burns KA, Vanden Heuvel JP. Modulation of PPAR activity via phosphorylation. *Biochim Biophys Acta* 2007;1771:952–60.
- Yamazaki K, Shimizu M, Okuno M, et al. Synergistic effects of RXR α and PPAR γ ligands to inhibit growth in human colon cancer cells-phosphorylated RXR α is a critical target for colon cancer management. *Gut* 2007;56:1557–63.
- Yu C, Markan K, Temple KA, Deplewski D, Brady MJ, Cohen RN. The nuclear receptor corepressors NCoR and SMRT decrease peroxisome proliferator-activated receptor γ transcriptional activity and repress 3T3-1 adipogenesis. *J Biol Chem* 2005;280:13600–5.
- Rosen ED, Spiegelman BM. Molecular regulation of adipogenesis. *Annu Rev Cell Dev Biol* 2000;16:145–71.
- Rosen ED, Walkey CJ, Puigserver P, Spiegelman BM. Transcriptional regulation of adipogenesis. *Genes Dev* 2000;14:1293–307.
- Spiegelman BM. PPAR- γ : adipogenic regulator and thiazolidinedione receptor. *Diabetes* 1998;47:507–14.
- Sertznig P, Seifert M, Tilgen W, Reichrath J. Present concepts and future outlook: function of peroxisome proliferator-activated receptors (PPARs) for pathogenesis, progression, and therapy of cancer. *J Cell Physiol* 2007;212:1–12.
- Burstein HJ, Demetri GD, Mueller E, Sarraf P, Spiegelman BM, Winer EP. Use of the peroxisome proliferator-activated receptor (PPAR) γ ligand troglitazone as treatment for refractory breast cancer: a phase II study. *Breast Cancer Res Treat* 2003;79:391–7.
- Smith MR, Manola J, Kaufman DS, et al. Rosiglitazone versus placebo for men with prostate carcinoma and a rising serum prostate-specific antigen level after radical prostatectomy and/or radiation therapy. *Cancer* 2004;101:1569–74.
- Demetri GD, Fletcher CD, Mueller E, et al. Induction of solid tumor differentiation by the peroxisome proliferator-activated receptor- γ ligand troglitazone in patients with liposarcoma. *Proc Natl Acad Sci U S A* 1999;96:3951–6.
- Green S, Tugwood JD, Issemann I. The molecular mechanism of peroxisome proliferator action: a model for species differences and mechanistic risk assessment. *Toxicol Lett* 1992;64–5 Spec No:131–9.
- Kroll TG, Sarraf P, Pecciarini L, et al. PAX8-1 fusion oncogene in human thyroid carcinoma [corrected]. *Science* 2000;289:1357–60.
- Cheung L, Messina M, Gill A, et al. Detection of the PAX8-PPAR γ fusion oncogene in both follicular thyroid carcinomas and adenomas. *J Clin Endocrinol Metab* 2003;88:354–7.
- Au AY, McBride C, Wilhelm KG, Jr., et al. PAX8-peroxisome proliferator-activated receptor γ (PPAR γ) disrupts normal PAX8 or PPAR γ transcriptional function and stimulates follicular thyroid cell growth. *Endocrinology* 2006;147:367–76.
- Araki O, Ying H, Furuya F, Zhu X, Cheng SY. Thyroid hormone receptor β mutants: dominant negative regulators of peroxisome proliferator-activated receptor γ action. *Proc Natl Acad Sci U S A* 2005;102:16251–6.
- Gupta RA, Tan J, Krause WF, et al. Prostacyclin-mediated activation of peroxisome proliferator-activated receptor δ in colorectal cancer. *Proc Natl Acad Sci U S A* 2000;97:13275–80.
- He TC, Chan TA, Vogelstein B, Kinzler KW. PPAR δ is an APC-regulated target of nonsteroidal anti-inflammatory drugs. *Cell* 1999;99:335–45.
- Saez E, Tontonoz P, Nelson MC, et al. Activators of the nuclear receptor PPAR γ enhance colon polyp formation. *Nat Med* 1998;4:1058–61.

39. Lefebvre M, Paulweber B, Fajas L, et al. Peroxisome proliferator-activated receptor γ is induced during differentiation of colon epithelium cells. *J Endocrinol* 1999;162:331–40.
40. Yang K, Fan KH, Lamprecht SA, et al. Peroxisome proliferator-activated receptor γ agonist troglitazone induces colon tumors in normal C57BL/6J mice and enhances colonic carcinogenesis in Apc^{1638 N/+} Mlh1^{+/-} double mutant mice. *Int J Cancer* 2005;116:495–9.
41. Linsalata M, Giannini R, Notarnicola M, Cavallini A. Peroxisome proliferator-activated receptor γ and spermidine/spermine N1-acetyltransferase gene expressions are significantly correlated in human colorectal cancer. *BMC Cancer* 2006;6:191.
42. Ikeda I, Tomimoto A, Wada K, et al. 5-Aminosalicylic acid given in the remission stage of colitis suppresses colitis-associated cancer in a mouse colitis model. *Clin Cancer Res* 2007;13:6527–31.
43. Rousseaux C, Lefebvre B, Dubuquoy L, et al. Intestinal antiinflammatory effect of 5-aminosalicylic acid is dependent on peroxisome proliferator-activated receptor- γ . *J Exp Med* 2005;201:1205–15.
44. Kohno H, Yasui Y, Suzuki R, Hosokawa M, Miyashita K, Tanaka T. Dietary seed oil rich in conjugated linolenic acid from bitter melon inhibits azoxymethane-induced rat colon carcinogenesis through elevation of colonic PPAR γ expression and alteration of lipid composition. *Int J Cancer* 2004;110:896–901.
45. Yasui Y, Suzuki R, Kohno H, et al. 9*trans*,11*trans* conjugated linoleic acid inhibits the development of azoxymethane-induced colonic aberrant crypt foci in rats. *Nutr Cancer* 2007;59:82–91.
46. Yasui Y, Hosokawa M, Kohno H, Tanaka T, Miyashita K. Troglitazone and 9*cis*,11*trans*,13*trans*-conjugated linolenic acid: comparison of their antiproliferative and apoptosis-inducing effects on different colon cancer cell lines. *Chemotherapy* 2006;52:220–5.
47. Sasaki T, Yoshida K, Shimura H, et al. Inhibitory effect of linoleic acid on transformation of IEC6 intestinal cells by *in vitro* azoxymethane treatment. *Int J Cancer* 2006;118:593–9.
48. Pandhare J, Cooper SK, Phang JM. Proline oxidase, a proapoptotic gene, is induced by troglitazone: evidence for both peroxisome proliferator-activated receptor γ -dependent and γ -independent mechanisms. *J Biol Chem* 2006;281:2044–52.
49. Park EY, Cho IJ, Kim SG. Transactivation of the PPAR-responsive enhancer module in chemopreventive glutathione S-transferase gene by the peroxisome proliferator-activated receptor- γ and retinoid X receptor heterodimer. *Cancer Res* 2004;64:3701–13.
50. Lu J, Imamura K, Nomura S, et al. Chemopreventive effect of peroxisome proliferator-activated receptor γ on gastric carcinogenesis in mice. *Cancer Res* 2005;65:4769–74.
51. Mehta RG, Williamson E, Patel MK, Koeffler HP. A ligand of peroxisome proliferator-activated receptor γ , retinoids, and prevention of preneoplastic mammary lesions. *J Natl Cancer Inst* 2000;92:418–23.
52. Suh N, Wang Y, Williams CR, et al. A new ligand for the peroxisome proliferator-activated receptor- γ (PPAR- γ), GW7845, inhibits rat mammary carcinogenesis. *Cancer Res* 1999;59:5671–3.
53. Yin Y, Russell RG, Dettin LE, et al. Peroxisome proliferator-activated receptor δ and γ agonists differentially alter tumor differentiation and progression during mammary carcinogenesis. *Cancer Res* 2005;65:3950–7.
54. Badawi AF, Eldeen MB, Liu Y, Ross EA, Badr MZ. Inhibition of rat mammary gland carcinogenesis by simultaneous targeting of cyclooxygenase-2 and peroxisome proliferator-activated receptor γ . *Cancer Res* 2004;64:1181–9.
55. Vogel U, Christensen J, Nexø BA, Wallin H, Friis S, Tjønneland A. Peroxisome proliferator-activated receptor- γ 2 Pro¹²Ala, interaction with alcohol intake and NSAID use, in relation to risk of breast cancer in a prospective study of Danes. *Carcinogenesis* 2007;28:427–34.
56. He G, Thuillier P, Fischer SM. Troglitazone inhibits cyclin D1 expression and cell cycling independently of PPAR γ in normal mouse skin keratinocytes. *J Invest Dermatol* 2004;123:1110–9.
57. He G, Muga S, Thuillier P, Lubet RA, Fischer SM. The effect of PPAR γ ligands on UV- or chemically-induced carcinogenesis in mouse skin. *Mol Carcinog* 2005;43:198–206.
58. Kim DJ, Murray IA, Burns AM, Gonzalez FJ, Perdew GH, Peters JM. Peroxisome proliferator-activated receptor- β/δ inhibits epidermal cell proliferation by down-regulation of kinase activity. *J Biol Chem* 2005;280:9519–27.
59. Yoshida K, Hirose Y, Tanaka T, et al. Inhibitory effects of troglitazone, a peroxisome proliferator-activated receptor γ ligand, in rat tongue carcinogenesis initiated with 4-nitroquinoline 1-oxide. *Cancer Sci* 2003;94:365–71.
60. Suzuki R, Kohno H, Suzui M, et al. An animal model for the rapid induction of tongue neoplasms in human c-Ha-ras proto-oncogene transgenic rats by 4-nitroquinoline 1-oxide: its potential use for preclinical chemoprevention studies. *Carcinogenesis* 2006;27:619–30.
61. Govindarajan R, Ratnasinghe L, Simmons DL, et al. Thiazolidinediones and the risk of lung, prostate, and colon cancer in patients with diabetes. *J Clin Oncol* 2007;25:1476–81.
62. Chang TH, Szabo E. Induction of differentiation and apoptosis by ligands of peroxisome proliferator-activated receptor γ in non-small cell lung cancer. *Cancer Res* 2000;60:1129–38.
63. Varley CL, Stahlschmidt J, Smith B, Stower M, Southgate J. Activation of peroxisome proliferator-activated receptor- γ reverses squamous metaplasia and induces transitional differentiation in normal human urothelial cells. *Am J Pathol* 2004;164:1789–98.
64. Varley CL, Stahlschmidt J, Lee WC, et al. Role of PPAR γ and EGFR signalling in the urothelial terminal differentiation programme. *J Cell Sci* 2004;117:2029–36.
65. Dominick MA, White MR, Sanderson TP, et al. Urothelial carcinogenesis in the urinary bladder of male rats treated with muraglitazar, a PPAR α/γ agonist: evidence for urolithiasis as the inciting event in the mode of action. *Toxicol Pathol* 2006;34:903–20.
66. Nissen SE, Wolski K. Effect of rosiglitazone on the risk of myocardial infarction and death from cardiovascular causes. *N Engl J Med* 2007;356:2457–71.
67. Dormandy JA, Charbonnel B, Eckland DJ, et al. Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (Prospective Pioglitazone Clinical Trial in Macrovascular Events): a randomised controlled trial. *Lancet* 2005;366:1279–89.
68. Charbonnel B, Dormandy J, Erdmann E, Massi-Benedetti M, Skene A. The prospective pioglitazone clinical trial in macrovascular events (PROactive): can pioglitazone reduce cardiovascular events in diabetes? Study design and baseline characteristics of 5238 patients. *Diabetes Care* 2004;27:1647–53.
69. Yadav N, Cheng D, Richard S, et al. CARM1 promotes adipocyte differentiation by coactivating PPAR γ . *EMBO Rep* 2008;9:193–8.
70. Fu M, Sun T, Bookout AL, et al. A nuclear receptor atlas: 3T3-1 adipogenesis. *Mol Endocrinol* 2005;19:2437–50.
71. Forman BM, Tontonoz P, Chen J, Brun RP, Spiegelman BM, Evans RM. 15-Deoxy- $\Delta^{12,14}$ -prostaglandin J₂ is a ligand for the adipocyte determination factor PPAR γ . *Cell* 1995;83:803–12.
72. Panigrahy D, Huang S, Kieran MW, Kaipainen A. PPAR γ as a therapeutic target for tumor angiogenesis and metastasis. *Cancer Biol Ther* 2005;4:687–93.
73. Plourde PV, Scott M. Duration of tamoxifen therapy and occurrence of endometrial cancer. *J Clin Oncol* 1995;13:2142.